

U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

PB-279 667

Distribution of Phytoplankton in Kentucky Lakes

Environmental Monitoring and Support Lab, Las Vegas, Nev

Jan 78

EPA-600/3-78-013
January 1978

PB 279 667

Ecological Research Series



REPRODUCED BY
**NATIONAL TECHNICAL
INFORMATION SERVICE**
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The nine series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy-Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ECOLOGICAL RESEARCH series. This series describes research on the effects of pollution on humans, plant and animal species, and materials. Problems are assessed for their long- and short-term influences. Investigations include formation, transport, and pathway studies to determine the fate of pollutants and their effects. This work provides the technical basis for setting standards to minimize undesirable changes in living organisms in the aquatic, terrestrial, and atmospheric environments.

TECHNICAL REPORT DATA (Please read instructions on the reverse before completing)		
1. REPORT NO. EPA-600/3-78-013	2.	3. REPORT ACCESSION NO. PB279667
4. TITLE AND SUBTITLE DISTRIBUTION OF PHYTOPLANKTON IN KENTUCKY LAKES		5. REPORT DATE January 1978
		6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S) W.D. Taylor, F.A. Hiatt, S.C. Hern, J.W. Hilgert, V.W. Lambou, F.A. Morris, M.K. Morris, L.R. Williams		8. PERFORMING ORGANIZATION REPORT NO.
9. PERFORMING ORGANIZATION NAME AND ADDRESS Environmental Monitoring and Support Laboratory Office of Research and Development U.S. Environmental Protection Agency Las Vegas, NV 89114		10. PROGRAM ELEMENT NO. 1BA608
		11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS U.S. Environmental Protection Agency-Las Vegas, NV Office of Research and Development Environmental Monitoring and Support Laboratory Las Vegas, NV 89114		13. TYPE OF REPORT AND PERIOD COVERED 03-07-73 to 11-14-73
		14. SPONSORING AGENCY CODE EPA/600/7
15. SUPPLEMENTARY NOTES Previously released in limited distribution as No. 683 in the Working Paper Series for the National Eutrophication Survey.		
16. ABSTRACT This is a data report presenting the species and abundance of phytoplankton in the 5 lakes sampled by the National Eutrophication Survey in the State of Kentucky. Results from the calculation of several water quality indices are also included (Nygaard's Trophic State Index, Palmer's Organic Pollution Index, and species diversity and abundance indices).		
17. KEY WORDS AND DOCUMENT ANALYSIS		
a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
*aquatic microbiology lakes *phytoplankton water quality	Kentucky lake eutrophication Nygaard's trophic indices Palmer's organic pollution indices Species diversity and abundance indices	06 C, M 08 H 13 B
18. DISTRIBUTION STATEMENT RELEASE TO PUBLIC	19. SECURITY CLASS (This Report) UNCLASSIFIED	21. NO. OF PAGES 36
	20. SECURITY CLASS (This page) UNCLASSIFIED	22. PRICE PC A03 / MF A01

EPA-600/3-78-013
January 1978

DISTRIBUTION OF PHYTOPLANKTON IN KENTUCKY LAKES

by

W. D. Taylor, F. A. Hiatt*, S. C. Hern, J. W. Hilgert*,
V. W. Lambou, F. A. Morris*, M. K. Morris*,
and L. R. Williams

Monitoring Operations Division
Environmental Monitoring and Support Laboratory
Las Vegas, Nevada 89114

*Department of Biological Sciences
The University of Nevada, Las Vegas
Las Vegas, Nevada 89154

ENVIRONMENTAL MONITORING AND SUPPORT LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
LAS VEGAS, NEVADA 89114

ENV-600-2-013
JANUARY 1980

DISTRIBUTION OF PHYTOPLANKTON IN KENTUCKY LAKE

DISCLAIMER

This report has been reviewed by the Environmental Monitoring and Support Laboratory-Las Vegas, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

Environmental Monitoring and Support Laboratory
U.S. Environmental Protection Agency
Las Vegas, Nevada 89160

Department of Biological Sciences
The University of Nevada, Las Vegas
Las Vegas, Nevada 89160


ENVIRONMENTAL MONITORING AND SUPPORT LABORATORY
OFFICE OF RESEARCH AND DEVELOPMENT
U.S. ENVIRONMENTAL PROTECTION AGENCY
LAS VEGAS, NEVADA 89160

FOREWORD

Protection of the environment requires effective regulatory actions which are based on sound technical and scientific information. This information must include the quantitative description and linking of pollutant sources, transport mechanisms, interactions, and resulting effects on man and his environment. Because of the complexities involved, assessment of specific pollutants in the environment requires a total systems approach which transcends the media of air, water, and land. The Environmental Monitoring and Support Laboratory-Las Vegas contributes to the formation and enhancement of a sound integrated monitoring data base through multidisciplinary, multimedia programs designed to:

- develop and optimize systems and strategies for monitoring pollutants and their impact on the environment
- demonstrate new monitoring systems and technologies by applying them to fulfill special monitoring needs of the Agency's operating programs

This report presents the species and abundance of phytoplankton in the 5 lakes sampled by the National Eutrophication Survey in the State of Kentucky, along with results from the calculation of several commonly used biological indices of water quality and community structure. These data can be used to biologically characterize the study lakes, and as baseline data for future investigations. This report was written for use by Federal, State, and local governmental agencies concerned with water quality analysis, monitoring, and/or regulation. Private industry and individuals similarly involved with the biological aspects of water quality will find the document useful. For further information contact the Water and Land Quality Branch, Monitoring Operations Division.



George B. Morgan
Director

Environmental Monitoring and Support Laboratory
Las Vegas

CONTENTS

	<u>Page</u>
Foreword	111
Introduction	1
Materials and Methods	2
Lakes and Site Selection	2
Sample Preparation	2
Examination	3
Quality Control	4
Results	5
Nygaard's Trophic State Indices	5
Palmer's Organic Pollution Indices	5
Species Diversity and Abundance Indices	7
Species Occurrence and Abundance	8
Literature Cited	9
Appendix. Summary of Phytoplankton Data	10

INTRODUCTION

The collection and analysis of phytoplankton data were included in the National Eutrophication Survey in an effort to determine relationships between algal characteristics and trophic status of individual lakes.

During spring, summer, and fall of 1973, the Survey sampled 250 lakes in 17 States. Over 700 algal species and varieties were identified and enumerated from the 743 water samples examined.

This report presents the species and abundance of phytoplankton in the 5 lakes sampled in the State of Kentucky (Table 1). The Nygaard's Trophic State (Nygaard 1949), Palmer's Organic Pollution (Palmer 1969), and species diversity and abundance indices are also included.

TABLE 1. LAKES SAMPLED IN THE STATE OF KENTUCKY

STORET #	LAKE NAME	COUNTY
2101	Lake Cumberland	Pulaski, McCreary, Russell, Wayne, Clinton
2102	Dale Hollow Reservoir	Cumberland, Clinton (Clay, Pickett, Overton in Tenn.)
2103	Herrington Lake	Boyle, Mercer, Garrard
2104	Kentucky Lake	Marshall, Lyon, Trigg, Livingston, Calloway (Henry, Stewart, Benton, Houston, Humphreys, Perry, Decatur in Tenn.)
2105	Barren River Reservoir	Allen, Barren

MATERIALS AND METHODS

LAKE AND SITE SELECTION

Lakes and reservoirs included in the Survey were selected through discussions with State water pollution agency personnel and U.S. Environmental Protection Agency Regional Offices (U.S. Environmental Protection Agency 1975). Screening and selection strongly emphasized lakes with actual or potential accelerated eutrophication problems. As a result, the selection was limited to lakes:

- (1) impacted by one or more municipal sewage treatment plant outfalls either directly into the lake or by discharge to an inlet tributary within approximately 40 kilometers of the lake;
- (2) 40 hectares or larger in size; and
- (3) with a mean hydraulic retention time of at least 30 days.

Specific selection criteria were waived for some lakes of particular State interest.

Sampling sites for a lake were selected based on available information on lake morphometry, potential major sources of nutrient input, and on-site judgment of the field limnologist (U.S. Environmental Protection Agency 1975). Primary sampling sites were chosen to reflect the deepest portion of each major basin in a test lake. Where many basins were present, selection was guided by nutrient source information on hand. At each sampling site, a depth-integrated phytoplankton sample was taken. Depth-integrated samples were uniform mixtures of water from the surface to a depth of 15 feet (4.6 meters) or from the surface to the lower limit of the photic zone representing 1 percent of the incident light, whichever was greater. If the depth at the sampling site was less than 15 feet (4.6 meters), the sample was taken from just off the bottom to the surface. Normally, a lake was sampled three times in 1 year, providing information on spring, summer, and fall conditions.

SAMPLE PREPARATION

Four milliliters (ml) of Acid-Lugol's solution (Prescott 1970) were added to each 130-ml sample from each site at the time of collection for preservation. The samples were shipped to the Environmental Monitoring and Support Laboratory, Las Vegas, Nevada, where equal volumes from each site were mixed to form two 130-ml composite samples for a given lake. One composite sample was put into storage and the other was used for the examination.

Prior to examination, the composite samples were concentrated by the settling method. Solids were allowed to settle for at least 24 hours prior to siphoning off the supernate. The volume of the removed supernate and the volume of the remaining concentrate were measured and concentrations determined. A small (8 ml) library subsample of the concentrate was then taken. The remaining concentrate was gently agitated to resuspend the plankton and poured into a capped, graduated test tube. If a preliminary examination of a sample indicated the need for a more concentrated sample, the contents of the test tube were further concentrated by repeating the settling method. Final concentrations varied from 15 to 40 times the original.

Permanent slides were prepared from concentrated samples after analysis was complete. A drop of superconcentrate from the bottom of the test tube was placed in a ring of clear Karo® Corn Syrup with phenol (a few crystals of phenol were added to each 100 ml of syrup) on a glass slide, thoroughly mixed, and topped with a coverglass. After the syrup at the edges of the coverglass had hardened, the excess was scraped away and the mount was sealed with clear fingernail polish. Permanent diatom slides were prepared by drying sample material on a coverglass, heating in a muffle furnace at 400° C for 45 minutes, and mounting in Hyrax®. Finally, the mounts were sealed with clear fingernail polish.

Backup samples, library samples, permanent sample slides, and Hyrax® mounted diatom slides are being stored and maintained at the U.S. Environmental Monitoring and Support Laboratory-Las Vegas.

EXAMINATION

The phytoplankton samples were examined with the aid of binocular compound microscopes. A preliminary examination was performed to precisely identify and list all forms encountered. The length of this examination varied depending on the complexity of the sample. An attempt was made to find and identify all of the forms present in each sample. Often forms were observed which could not be identified to species or to genus. Abbreviated descriptions were used to keep a record of these forms (e.g., lunate cell, blue-green filament, Navicula #1). Diatom slides were examined using a standard light microscope. If greater resolution was essential to accurately identify the diatoms, a phase-contrast microscope was used.

After the species list was compiled, phytoplankton were enumerated using a Neubauer Counting Chamber with a 40X objective lens and a 10X ocular lens. All forms within each field were counted. The count was continued until a minimum of 100 fields had been viewed, or until the dominant form had been observed a minimum of 100 times.

®Registered Trademark

QUALITY CONTROL

Internal quality control checks on species identifications and counts were performed on a regular basis between project phycologists at the rate of 7 percent. Although an individual had primary responsibility for analyzing a sample, taxonomic problems were discussed among the phycologists.

Additional quality control checks were performed on the Survey samples by Dr. G. W. Prescott of the University of Montana at the rate of 5 percent. Quality control checks were made on 75 percent of these samples to verify species identifications while checks were made on the remaining 25 percent of the samples to verify genus counts. Presently, the agreement between quality control checks for species identification and genus enumerations is satisfactory.

RESULTS

The Appendix summarizes all of the phytoplankton data collected from the State by the Survey. It is organized by lake, including an alphabetical phytoplankton species list with concentrations for individual species given by sampling date. Results from the application of several indices are presented (Nygaard's Trophic State, Palmer's Organic Pollution, and species diversity and abundance). Each lake has been assigned a four digit STORET number. [STORET (STOrage and RETrieval) is the U.S. Environmental Protection Agency's computer system which processes and maintains water quality data.] The first two digits of the STORET number identify the State; the last two digits identify the lake.

NYGAARD'S TROPHIC STATE INDICES

Five indices devised by Nygaard (1949) were proposed under the assumption that certain algal groups are indicative of levels of nutrient enrichment. These indices were calculated in order to aid in determining the surveyed lakes' trophic status. As a general rule, Cyanophyta, Euglenophyta, centric diatoms, and members of the Chlorococcales are found in waters that are eutrophic (rich in nutrients), while desmids and many pennate diatoms generally cannot tolerate high nutrient levels and so are found in oligotrophic waters (poor in nutrients).

In applying the indices to the Survey data, the number of taxa in each major group was determined from the species list for each sample. The ratios of these groups give numerical values which can be used as a biological index of water richness. The five indices and the ranges of values established for Danish lakes by Nygaard for each trophic state are presented in Table 2. The appropriate symbol, (E) eutrophic and (O) oligotrophic, follows each calculated value in the tables in the Appendix. A question mark (?) was entered in these tables when the calculated value was within the range of both classifications.

PALMER'S ORGANIC POLLUTION INDICES

Palmer (1969) analyzed reports from 165 authors and developed algal pollution indices for use in rating water samples with high organic pollution. Two lists of organic pollution-tolerant forms were prepared, one containing 20 genera, the other, 20 species (Tables 3 and 4). Each form was assigned a pollution index number ranging from 1 for moderately tolerant forms to 6 for extremely tolerant forms. Palmer based the index numbers on occurrence records and/or where emphasized by the authors as being especially tolerant of organic pollution.

TABLE 2. NYGAARD'S TROPHIC STATE INDICES ADAPTED FROM HUTCHINSON (1967)

Index	Calculation	Oligotrophic	Eutrophic
Myxophycean	<u>Myxophyceae</u> <u>Desmidiaceae</u>	0.0-0.4	0.1-3.0
Chlorophycean	<u>Chlorococcales</u> <u>Desmidiaceae</u>	0.0-0.7	0.2-9.0
Diatom	<u>Centric Diatoms</u> <u>Pennate Diatoms</u>	0.0-0.3	0.0-1.75
Euglenophyte	<u>Euglenophyta</u> <u>Myxophyceae + Chlorococcales</u>	0.0-0.3	0.0-1.0
Compound	<u>Myxophyceae + Chlorococcales +</u> <u>Centric Diatoms + Euglenophyta</u> <u>Desmidiaceae</u>	0.0-1.0	1.2-25

TABLE 3. ALGAL GENUS POLLUTION INDEX
(Palmer 1969)

Genus	Pollution Index
<i>Anacystis</i>	1
<i>Ankistrodesmus</i>	2
<i>Chlamydomonas</i>	4
<i>Chlorella</i>	3
<i>Closterium</i>	1
<i>Cyclotella</i>	1
<i>Euglena</i>	5
<i>Gomphonema</i>	1
<i>Lepocinclis</i>	1
<i>Melosira</i>	1
<i>Micractinium</i>	1
<i>Navicula</i>	3
<i>Nitzschia</i>	3
<i>Oscillatoria</i>	5
<i>Pandorina</i>	1
<i>Phacus</i>	2
<i>Phormidium</i>	1
<i>Scenedesmus</i>	4
<i>Stigeoclonium</i>	2
<i>Synedra</i>	2

TABLE 4. ALGAL SPECIES POLLUTION INDEX (Palmer 1969)

Species	Pollution Index
<i>Ankistrodesmus falcatus</i>	3
<i>Arthrospira jenneri</i>	2
<i>Chlorella vulgaris</i>	2
<i>Cyclotella meneghiniana</i>	2
<i>Euglena gracilis</i>	1
<i>Euglena viridis</i>	6
<i>Gomphonema parvulum</i>	1
<i>Melosira varians</i>	2
<i>Navicula cryptocephala</i>	1
<i>Nitzschia acicularis</i>	1
<i>Nitzschia palea</i>	5
<i>Oscillatoria chlorina</i>	2
<i>Oscillatoria limosa</i>	4
<i>Oscillatoria princeps</i>	1
<i>Oscillatoria putrida</i>	1
<i>Oscillatoria tenuis</i>	4
<i>Pandorina morum</i>	3
<i>Scenedesmus quadricauda</i>	4
<i>Stigeoclonium tenue</i>	3
<i>Synedra ulna</i>	3

In analyzing a water sample, any of the 20 genera or species of algae present in concentrations of 50 per milliliter or more are recorded. The pollution index numbers of the algae present are totaled, providing a genus score and a species score. Palmer determined that a score of 20 or more for either index can be taken as evidence of high organic pollution, while a score of 15 to 19 is taken as probable evidence of high organic pollution. Lower figures suggest that the organic pollution of the sample is not high, that the sample is not representative, or that some substance or factor interfering with algal persistence is present and active.

SPECIES DIVERSITY AND ABUNDANCE INDICES

"Information content" of biological samples is being used commonly by biologists as a measure of diversity. Diversity in this connection means the degree of uncertainty attached to the specific identity of any randomly selected individual. The greater the number of taxa and the more equal their proportions, the greater the uncertainty, and hence, the diversity (Pielou 1966). There are several methods of measuring diversity, e.g., the formulas given by Brillouin (1962) and Shannon and Weaver (1963). The method which is appropriate depends on the type of biological sample on hand.

Pielou (1966) classifies the types of biological samples and gives the measure of diversity appropriate for each type. The Survey phytoplankton samples are what she classifies as larger samples (collections in Pielou's terminology) from which random subsamples can be drawn. According to Pielou, the average diversity per individual for these types of samples can be estimated from the Shannon-Wiener formula (Shannon and Weaver 1963):

$$H = -\sum_{i=1}^S P_i \log_x P_i$$

where P is the proportion of the i th taxon in the sample, which is calculated from n_i/N ; n_i is the number of individuals per milliliter of the i th taxon; N is the total number of individuals per ml; and S is the total number of taxa.

However, Basharin (1959) and Pielou (1966) have pointed out that H calculated from the subsample is a biased estimator of the sample H , and if this bias is to be accounted for, we must know the total number of taxa present in the sample since the magnitude of this bias depends on it.

Pielou (1966) suggests that if the number of taxa in the subsample falls only slightly short of the number in the larger sample, no appreciable error will result in considering S , estimated from the subsample, as being equal to the sample value. Even though considerable effort was made to find and identify all taxa, the Survey samples undoubtedly contain a fair number of rare phytoplankton taxa which were not encountered.

In the Shannon-Wiener formula, an increase in the number of taxa and/or an increase in the evenness of the distribution of individuals among taxa will increase the average diversity per individual from its minimal value of zero. Sager and Hasler (1969) found that the richness of taxa was of minor importance in determination of average diversity per individual for phytoplankton and they concluded that phytoplankton taxa in excess of the 10 to 15 most abundant ones have little effect on H , which was verified by our own calculations. Our counts are in number per milliliter and since logarithms to the base 2 were used in our calculations, H is expressed in units of bits per individual. When individuals of a taxon were so rare that they were not counted, a value of 1/130 per milliliter or 0.008 per milliliter was used in the calculations since at least one individual of the taxon must have been present in the collection.

A Survey sample for a given lake represents a composite of all phytoplankton collected at different sampling sites on a lake during a given sampling period. Since the number of samples (M) making up a composite is a function of both the complexity of the lake sampled and its size, it should affect the richness-of-taxa component of the diversity of our phytoplankton collections. The maximum diversity ($\text{Max}H$) (i.e., when the individuals are distributed among the taxa as evenly as possible) was estimated from $\log_2 S$, the total diversity (D) was calculated from H/N , and the evenness component of diversity (J) was estimated from $H/\text{Max}H$ (Pielou 1966). Also given in the Appendix are L (the mean number of individuals per taxa per milliliter) and K (the number of individuals per milliliter of the most abundant taxon in the sample).

Zand (1976) suggests that diversity indices be expressed in units of "sits", i.e., in logarithms to base S (where S is the total number of taxa in the sample) instead of in "bits", i.e., in logarithms to base 2. Zand points out that the diversity index in sits per individual is a normalized number ranging from 1 for the most evenly distributed samples to 0 for the least evenly distributed samples. Also, it can be used to compare different samples, independent of the number of taxa in each. The diversity in bits per individual should not be used in direct comparisons involving various samples which have different numbers of species. Since $\text{Max}H$ equals $\log_2 S$, the expression in sits is equal to $\log_S S$, or 1. Therefore diversity in sits per individual is numerically equivalent to J , the evenness component for the Shannon-Wiener formula.

SPECIES OCCURRENCE AND ABUNDANCE

The alphabetic phytoplankton species list for each lake, presented in the Appendix, gives the concentrations of individual species by sampling date. Concentrations are in cells, colonies, or filaments (CEL, COL, FIL) per milliliter. An "X" after a species name indicates the presence of the species on that date in such a low concentration that it did not show up in the count. A blank space indicates that the organism was not found in the sample collected on that date. Column S is used to designate the examiner's subjective opinion of the five dominant taxa in a sample, based upon relative size and concentration of the organism. The percent column (%) presents, by abundance, the percentage composition of each taxon.

LITERATURE CITED

- Basharin, G. P. 1959. On a statistical estimate for the entropy of a sequence of independent random variables, pp. 333-336. In N. Artin (ed.), Theory of Probability and Its Applications (translation of "Teoriya Veroyatnosc i ee Premeneniya") 4. Society for Industrial and Applied Mathematics, Philadelphia.
- Brillouin, L. 1962. Science and Information Theory (2nd ed.). Academic Press, New York. 351 pp.
- Hutchinson, G. E. 1967. A Treatise on Limnology. II. Introduction to Lake Biology and the Limnoplankton. John Wiley and Sons, Inc., New York. 1,115 pp.
- Nygaard, G. 1949. Hydrobiological studies of some Danish ponds and lakes. II. (K danske Vidensk. Selsk.) Biol. Sci. 7:293.
- Palmer, C. M. 1969. A composite rating of algae tolerating organic pollution. J. Phycol. 5:78-82.
- Pielou, E. C. 1966. The measurement of diversity in different types of biological collections. J. Theor. Biol. 13:131-144.
- Prescott, G. W. 1970. How to Know the Freshwater Algae. William C. Brown Company, Dubuque. 348 pp.
- Sager, P. E. and A.D. Hasler. 1969. Species diversity in lacustrine phytoplankton. I. The components of the index of diversity from Shannon's formula. Amer. Natur. 103(929):51-59.
- Shannon, C. E. and W. Weaver. 1963. The Mathematical Theory of Communication. University of Illinois Press, Urbana. 117 pp.
- U.S. Environmental Protection Agency. 1975. National Eutrophication Survey Methods 1973-1976. Working Paper No. 175. Environmental Monitoring and Support Laboratory. Las Vegas, Nevada, and Corvallis Environmental Research Laboratory, Corvallis, Oregon. 91 pp.
- Zand, S. M. 1976. Indexes associated with information theory in water quality. Journal WPCF. 48(8):2026-2031.

APPENDIX. SUMMARY OF PHYTOPLANKTON DATA

This appendix was generated by computer. Because it was only possible to use upper case letters in the printout, all scientific names are printed in upper case and are not italicized.

The alphabetic phytoplankton lists include taxa without species names (e.g., EUNOTIA, EUNOTIA #1, FLAGELLATE, FLAGELLATES, MICROSYSTIS INCERTA ?, CHLOROPHYTAN COCCOID CELLED COLONY). When species determinations were not possible, symbols or descriptive phrases were used to separate taxa for enumeration purposes. Each name on a list, however, represents a unique species different from any other name on the same list, unless otherwise noted, for counting purposes.

Numbers were used to separate unidentified species of the same genus. A generic name listed alone is also a unique species. A question mark (?) is placed immediately after the portion of a name which was assigned with uncertainty. Numbered, questioned, or otherwise designated taxa were established on a lake-by-lake basis; therefore NAVICULA #2 from lake A cannot be compared to NAVICULA #2 from lake B. Pluralized categories (e.g., FLAGELLATES, CENTRIC DIATOMS, SPP.) were used for counting purposes when taxa could not be properly differentiated on the counting chamber.

LAKE NAME: LAKE CUMBERLAND
STORY NUMBER: 2101

NYGAARD TROPIC STATE INDICES

DATE	05 29 73	08 21 73	10 25 73
MYXOPHYCEAN	02/0 F	3.00 E	03/0 E
CHLOROPHYCEAN	01/0 E	1.00 E	01/0 E
EUGLENOPHYTE	07/0 7	0.25 E	0.25 E
DIATOM	0.80 E	0.25 7	0.33 E
COMPOUND	07/0 E	6.00 E	07/0 E

PALMER'S ORGANIC POLLUTION INDICES

DATE	05 29 73	08 21 73	10 25 73
GENUS	00	03	03
SPECIES	00	00	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE	05 29 73	08 21 73	10 25 73
AVERAGE DIVERSITY	2.28	1.13	2.34
NUMBER OF TAXA	17.00	15.00	17.00
NUMBER OF SAMPLES COMPOSITED	7.00	7.00	7.00
MAXIMUM DIVERSITY	4.09	3.91	4.09
TOTAL DIVERSITY	2665.32	7133.69	2686.32
MEAN NUMBER OF INDIVIDUALS/TAXA	1169.00	6313.00	1148.00
EVENNESS COMPONENT	0.56	0.29	0.57
MEAN NUMBER OF INDIVIDUALS/TAXA	68.76	420.87	67.53
NUMBER/ML OF MOST ABUNDANT TAXON	447.00	5132.00	628.00

LAKE NAME: LAKE CUMBERLAND
STREET NUMBER: 2101

CONTINUED

TAXA	FORM	05 29 73			08 21 73			10 25 73		
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
ANABAENA	FIL			X						
ANABAENA #1	FIL					0.3	19			X
ANABAENA #2	FIL					1.81.3	5132			X
ANCHOCNEIS	CEL					0.3	19			628
APHANOTHECE	COL								1154.7	
ASTERICNELLA FORMOSA	CEL		1.1	13						
CERATIUM HIRUNDINELLA	CEL			X		0.3	19			
CCSMARIUM	CEL					3.9	248			
CRYPTOMONAS	CEL		6.8	79						
CYANOPHYTAN FILAMENT	CEL					0.3	19			
CYMBELLA	FIL					2.1	133		5.3	61
DINOBRYON	CEL									X
DINOBRYON DIVERGENS	CEL								1.3	15
DINOFAGELLATE #1	CEL		9.0	105						
DINOFAGELLATE #2	CEL								2.7	31
FLAGELLATES	CEL									
FRAGILARIA CROTCHENSIS	CEL	2138.2		447	2	9.1	572	3	9.3	107
MALLCMCNAS PSEUDOCORONATA	CEL	1133.7		394	5	0.9	57	5	8.0	92
MELOSIRA #2	CEL						X		4.3	15
MELOSIRA DISTANS	CEL			X					4.0	46
MELOSIRA VARIANS	CEL			X						
NAVICULA	CEL	2.2		26						
NAVICULA SALINARTUM	CEL			X			X			
V. INTERMEDIA	CEL									
NITZSCHIA	CEL			X						
OSCILLATORIA	CEL									X
OSCILLATORIA #1	FIL								1.3	15
PANDORINA MORUM	FIL		1.1	13						
PHACUS	COL			X						
SCENEDESMUS	CEL									
STEPHANODISCUS	COL	13	5.6	66		0.3	19			X

THE CUMBERLAND
2101
CONTINUED

CONTINUED

SYNECRA #1
SYNECRA DELICATISSIMA
V. ANGUSTISSIMA ?
TETRAEDRON MINIMUM
TRACHELOMONAS

1159

LAKE NAME: DALE HOLLOW RES.
SECRET NUMBER: 2102

NYGAARD TROPHIC STATE INDICES

DATE	05	18	73	08	18	73	10	24	73
MYXOPHYCEAN	04/0	E	0/0	E	0/0	0	01/0	E	
CHLOROPHYCEAN	01/0	E	02/0	E	02/0	E	04/0	E	
EUGLENOPHYCEAN	0-20	7	0/02	7	0/02	7	0/05	7	
DIATOM	0-40	E	1-00	E	1-00	E	0/05	7	
COMPOUND	08/0	E	06/0	E	06/0	E	05/0	E	

PALMER'S ORGANIC POLLUTION INDICES

DATE	05	18	73	08	18	73	10	24	73
GENUS	00		00		01		00		00
SPECIES	00		00		00		00		00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE	05	18	73	08	18	73	10	24	73
AVERAGE DIVERSITY	H	2-12		2-28		2-13			
NUMBER OF TAXA	S	16-00		17-00		14-00			
NUMBER OF SAMPLES COMPOSITED	M	6-00		6-00		6-00			
MAXIMUM DIVERSITY	MAXH	4-00		4-09		3-81			
TOTAL DIVERSITY	D	493-86		2948-04		1433-49			
TOTAL NUMBER OF INDIVIDUALS/ML	N	233-00		1293-00		673-00			
EVENNESS COMPONENT	J	0-53		0-56		0-56			
MEAN NUMBER OF INDIVIDUALS/TAXA	L	14-56		76-06		48-07			
NUMBER/ML OF MOST ABUNDANT TAXON	K	112-00		617-00		412-00			

CONTINUED

15

LAKE NAME: DALE P-OLLOH RES.
SYCRET NUMBER: 2102

CONTINUED

TAXA	FORM	05 18 73		08 18 73		10 24 73	
		SC	ALGAL UNITS PER ML	SC	ALGAL UNITS PER ML	SC	ALGAL UNITS PER ML
SCENEDESINUS #3	COL						
SYNEDRA #1	CEL			1.9	24	7.3	49
SYNEDRA #2	CEL			0.9	12		
TOTAL			233		1293		673

LAKE NAME: HERRINGTON LAKE
SECRET NUMBER: 2103

NYGAARD TROPHIC STATE INDICES				
DATE	05 25 73	08 20 73	10 23 73	
MYXOPHYCEAN	04/0 E	10.0 E	03/0 E	
CHLOROPHYCEAN	05/0 E	3.00 E	08/0 E	
EUGLENOPHYCEAN	0/09 ?	0.08 ?	0.27 E	
DIATOM	0.60 E	0.25 ?	0.40 E	
COMPOUND	12/0 E	15.0 E	18/0 E	

PALMER'S ORGANIC POLLUTION INDICES

DATE	05 26 73	08 20 73	10 23 73
GENUS	11	14	10
SPECIES	00	04	00

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE	05 26 73	08 20 73	10 23 73
AVERAGE DIVERSITY	1.48	2.90	1.70
NUMBER OF TAXA	21.00	24.00	32.00
NUMBER OF SAMPLES COMPOSITED	4.00	4.00	4.00
MAXIMUM DIVERSITY	4.39	4.58	5.00
TOTAL DIVERSITY	15996.88	16173.30	17715.70
TOTAL NUMBER OF INDIVIDUALS/ML	10606.00	5577.00	10421.00
EVENNESS COMPONENT	0.34	0.63	0.34
MEAN NUMBER OF INDIVIDUALS/TAXA	505.05	232.38	325.66
NUMBER/ML OF MOST ABUNDANT TAXON	4970.00	1711.00	7501.00

LAKE NAME: HERRINGTON LAKE
SYRET NUMBER: 2103

CONTINUED

TAXA	FORM	05 26 73			08 20 73			10 23 73		
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
ACHNANTHES MICROCEPHALA ?	CEL									X
ACTINASTRUM HANTZSCHII	COL								1.7	175
AMPHICRA	CEL									X
ANABAENA	FIL		0.2	22						
ANABAENA #1	FIL					4.3	242			
ANABAENOPSIS	FIL					0.5	29			
ASTERIONELLA FORMOSA	CEL	14	0.8	89						
COELASTRUM MICROPORUM	COL			X						
COELASTRUM SPHAERICUM	COL					0.3	14			X
COSMARUM	CEL					0.3	14			
CYCLOTELLA	CEL		0.8	89						
CYCLOTELLA STELLIGERA	CEL							4	2.4	246
DACTYLOCOCCOPSIS	CEL		0.2	22		1.5	86		1.7	176
DINOFAGELLATE	CEL		0.2	22						
DINOFAGELLATE #1	CEL					0.8	43			X
DINOFAGELLATE #2	CEL						X			X
EUGLENA #1	CEL									70
EUGLENA #2	CEL					0.3	14		0.3	35
FLAGELLATES	CEL	11	46.9	4970		1126.9	1498	11	172.0	7501
FRAGILARIA CROTONEENSIS	CEL	21	46.9	4970						
GONIUM PECTORALE	CEL			X						
HANTZSCHIA	CEL									X
KIRCHNERIELLA	CEL								0.3	35
LYNGBYA LIMNETICA	FIL					0.5	29			
PELCSIPA #2	CEL	15	0.6	67						
MELOSIRA #4	CEL								1.4	141
PELOSIRA DISTANS	CEL							12	9.8	1021
MERISOMEDIA MARSSONII	COL							15	2.0	211
MERISOMEDIA TENUISSIMA	COL					1.3	71			
MESOSTIGMA	CEL								1.0	106
PICRCCYSTIS INCERTA	CEL					0.5	29			X
NAVICULA #1	CEL									X

LAKE NAME: HERRINGTON LAKE
STRET NUMBER: 2103

CONTINUED

TAXA	FORM	05 26 73			08 20 73			10 23 73		
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
NAVICULA #2	CEL			X						X
NITZSCHIA ? #2	CEL								0.7	70
NITZSCHIA #1	CEL									X
NITZSCHIA #3	CEL			X						
NITZSCHIA HOLSATICA ?	FIL			X		1.5	86			
OSCILLATORIA	FIL			67		210.5	585			
OSCILLATORIA LINNETICA	COL		0.6	X			X			
PANDORINA PCRUM	CEL									
PHACUS ALATUS ?	CEL									
RAPHIDIOPSIS	FIL					4130.7	1711		0.3	35
SCENEDESMUS #1	COL		0.8	89						X
SCENEDESMUS #2	COL		0.4	44					0.3	35
SCENEDESMUS #3	COL									X
SCENEDESMUS DIMORPHUS	COL		0.2	22						X
SCENEDESMUS QUADRICAUDA	COL					1.8	100			
STEPHANODISCUS	CEL	3	1.3	133						
STEPHANODISCUS ?	CEL					3111.5	642		5.4	563
SURIELLA	CEL									X
SYNEDRA #1	CEL									X
SYNEDRA #2	CEL					4.3	242			X
SYNEDRA #3	CEL									X
SYNEDRA DELICATISSIMA	CEL			X		1.0	57			
TETRAEDRON MINIMUM	CEL					0.3	14			
TETRAEDRON MUTICUM	CEL			X		1.3	71			X
TOTAL				10606			5577			10421

LAKE NAME: KENTUCKY LAKE
 STORER NUMBER: 2104

NYGAARD TROPIC STATE INDICES

DATE	05	16	73	08	13	73	10	20	73
MYXOPHYCEAN	4.00	E	1.33	E	1.43	E			
CHLOROPHYCEAN	8.00	E	2.67	E	1.57	E			
EUGLENOPHYTE	0.33	E	0.25	E	0.14	E			
DIATOM	1.00	E	1.20	E	1.17	E			
COMPOUND	24.0	E	6.00	E	4.43	E			

PALMER'S ORGANIC POLLUTION INDICES

DATE	05	16	73	08	13	73	10	20	73
GENUS			04		15			17	
SPECIES			00		02			01	

SPECIES DIVERSITY AND ABUNDANCE INDICES

DATE	05	16	73	08	13	73	10	20	73
AVERAGE DIVERSITY	H	2.37		3.64		3.42			
NUMBER OF TAXA	S	39.00		52.00		46.00			
NUMBER OF SAMPLES COMPOSITED	M	17.00		10.00		9.00			
MAXIMUM DIVERSITY	MAXH	5.29		5.70		5.52			
TOTAL DIVERSITY	D	13049.22		21752.64		26556.72			
TOTAL NUMBER OF INDIVIDUALS/ML	N	5506.00		5976.00		7766.00			
EVENNESS COMPONENT	J	0.45		0.64		0.62			
MEAN NUMBER CF INDIVIDUALS/TAXA	L	141.18		114.92		168.83			
NUMBER/ML OF MOST ABUNDANT TAXON	K	2225.00		1002.00		2943.00			

LAKE NAME: KENTUCKY LAKE
STORET NUMBER: 2104

CONTINUED

TAXA	FORM	05 16 73			08 13 73			10 20 73		
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
ACHNANTHES MICROCEPHALA ?	CEL									
ANABAENA #1	FIL									128
ANABAENA #2	FIL					1.5	88		1.6	512
ANABAENA #3	FIL								1.1	85
ANABAENOPSIS	FIL								0.6	43
ANABAENOPSIS PHILIPPINENSIS	FIL								0.6	43
ANKISTRODESCHMUS	CEL					5.4	324		0.6	43
APHANIZCUMEN	COL								0.6	43
APHANIZCUMEN ?	FIL					0.5	29			
ASTERICNELLA FORMOSA	CEL		0.8	42					2.2	171
CENTRITRACTUS ?	CEL			X						
CERATIUM HIRUNDINELLA	CEL			X						
CLCSTERIOPSIS ?	CEL									
CLCSTERIUM ? #1	CEL									
CLCSTERIUM ? #3	CEL									
CLCSTERIUM #2	CEL									
COELASTRUM RETICULATUM	CEL									
COELASTRUM SPHAERICUM	COL									X
CCSMARIUM	COL			X					0.6	X
CCSMARIUM #1	CEL									
CCSMARIUM #2	CEL									
CRUCIGENTIA APICULATA	CEL					0.5	29			
CRYPTOMONAS	COL									
CYCLOTHELLA MENEGHINIANA	CEL									
CYCLOTHELLA STELLIGERA	CEL									
CYMBELLA	CEL		0.4	20		3.0	177		0.6	43
DACTYLOCCOPIS	CEL		0.2	11					1.1	85
DICTYOSPHAERIUM PULCHELLUM	CEL		0.5	30						
DINOBRYON	CEL									
DINOBRYON EAVARICUM	CEL		0.1	6						
DINOFAGELLATE	CEL			X						
DINOFAGELLATES	CEL			X		0.5	29			

LAKE NAME: KENTUCKY LAKE
STOREY NUMBER: 2104

CONTINUED

TAXA	FORM	05 16 73				08 13 73				10 20 73			
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
EUASTRUM DENTICULATUM	COL												
EUASTRUM DENTICULATUM ?	COL												
EUGLENA	CEL			X			29						X
EUGLENA #1	CEL			X									
EUGLENA #2	CEL						29						
EUGLENA #3	CEL						X						X
FLAGELLATES	CEL						X						
GYROSTOMA	CEL	12	31.8	1793	12	15.3	913	3	11.0	953			
GYROSTOMA ?	CEL			X									
LYNGBYA #2	CEL												
LYNGBYA CONTORTA	FIL						X						
MALLONAS PSEUDOCORONATA ?	FIL						737						
MELOSIRA #2	CEL		1.0	56		12.3	29		0.6	43			43
MELOSIRA #3	CEL	13	9.1	503		0.5			0.6	43			
MELOSIRA #4	CEL												
MELOSIRA #5	CEL	11	40.4	2225	3	7.9	471	12	8.8	683			
MELOSIRA DISTANS	CEL				5	3.9	236						
MELOSIRA VARIANS	CEL				11	15.8	1002		4.4	341			128
MERISOPEDIA	CEL	5	4.5	250		5.4	324		1.6	85			
MERISOPEDIA MARSSONII	CEL		0.4	21					1.1				
MICRACTINIUM	COL		0.5	28									
MICROCYSTIS AERUGINOSA	COL								0.6	43			
NAVICULA	COL			X		0.5	29						
NITZSCHIA	CEL												
NITZSCHIA #2	CEL												
NITZSCHIA ACICULARIS	CEL			X		1.0	59		1.1	85			
OSCILLATORIA	CEL		1.3	73		0.5	29						
OSCILLATORIA LIMNETICA	FIL												
PANDORINA MORUM	FIL					1.5	88		4.4	341			
PEDIASTRUM BIRADIATUM	FIL			X			X	5	6.6	512			
V. LONGECORNUTUM	COL		0.1	6	4	14.8	884	1	37.9	2943			
PEDIASTRUM SIMPLEX	COL						X						
V. DUODENARIUM	COL						X						
	COL						X						X

LAKE NAME: KENTUCKY LAKE
STCRET NUMBER: 2104

CONTINUED

TAXA	FORM	05 16 73			08 13 73			10 20 73		
		IS	SC	ALGAL UNITS PER ML	S	SC	ALGAL UNITS PER ML	S	SC	ALGAL UNITS PER ML
PEDIASTRUM TETRAS	COL									
V. TETRAEDON	CEL		1.5	83			X			
PENNATE CIATOM	CEL						X			
PHACUS PYRUM	CEL			X						
PHACUS TERTUS	CEL									
RHIZOLENTIA	CEL			X						X
SCENEDESMUS	COL		0.3	14						
SCENEDESMUS #1	COL					1.0	59			X
SCENEDESMUS #2	COL					0.5	29			X
SCENEDESMUS ABUNCANS	COL			X						X
SCENEDESMUS BI JUGA	COL									
V. ALTERNANS	COL									
SCENEDESMUS DIMORPHUS	COL						X			X
SCENEDESMUS DISPAR	COL						X		1.1	85
SCENEDESMUS PROTUBERANS	COL						X			
SCENEDESMUS QUADRICAUDA	COL			X						
SCENEDESMUS QUADRICAUDA ?	COL					0.5	29		0.6	43
V. ALTERNANS	COL		0.1	6						
SCHROEDERIA SETIGERA	CEL					0.5	29			
STALRASTRUM #1	CEL									
STALRASTRUM #2	CEL						X			X
STALRASTRUM #3	CEL									X
STEPHANODISCUS	CEL		6.4	354		4.9	295		2.7	213
SURIELLA ANGUSTATA	CEL			X						
SYNEDRA	CEL		0.5	25						
SYNEDRA #1	CEL									
TETRAEDRON MINIMUM	CEL						X		0.6	X
V. SCROBICULATUM	CEL									43
TETRAEDRON TRIGONUM	CEL								0.6	43
V. GPACILE	CEL									
TETRASTRUM ELEGANS	COL			X			X			

LAKE NAMES: KENTUCKY LAKE
 STREET NUMBER: 2104

CONTINUED

TAXA	FORM	05 16 73			08 13 73			10 20 73		
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
TRACHELOMONAS #1	CEL									
TRACHELOMONAS #2	CEL			X			X			X
TOTAL				5506			5976			7766

LAKE NAME: BARREN RIVER RES.
STORET NUMBER: 2105

NYGAARD TRCPHC STATE INDICES									
DATE	05	18	73	08	11	73	10	23	73
MYXOPHYCEAN	02/0	E		04/0	E		2.00	E	
CHLOROPHYCEAN	07/0	E		01/0	E		4.47	E	
EUGLENCYPHYTE	0.11	?		0.80	E		0.20	?	
DIATOM	0.80	E		0.50	E		0.71	E	
COMPOUND	18/0	E		14/0	E		9.67	E	

PALMER'S ORGANIC POLLUTION INDICES									
DATE	05	18	73	08	11	73	10	23	73
GENUS	07			16			11		
SPECIES	00			00			04		

SPECIES DIVERSITY AND ABUNDANCE INDICES									
DATE	05	18	73	08	11	73	10	23	73
AVERAGE DIVERSITY	H	2.68		1.71			3.25		
NUMBER OF TAXA	S	34.00		25.00			45.00		
NUMBER OF SAMPLES COMPOSITED	M	11.00		11.00			10.00		
MAXIMUM DIVERSITY	MAXH	5.09		4.64			5.49		
TOTAL DIVERSITY	D	28043.52		24871.95			10959.00		
MEAN NUMBER OF INDIVIDUALS/ML	N	10464.00		14545.00			3372.00		
EVENNESS COMPONENT	J	0.53		0.37			0.59		
MEAN NUMBER OF INDIVIDUALS/TAXA	L	307.76		581.80			74.93		
NUMBER/ML OF MOST ABUNDANT TAXON	K	4402.00		9775.00			968.00		

LAKE NAME: GARDEN RIVER RES.
STORE NUMBER: 2105

CONTINUED

TAXA	FORM	05 10 73		08 11 73		10 23 73	
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
ACHNANTHES	CEL			X			15
ACHNANTHES LANCECLATA	CEL						
ACHNANTHES MICROCEPHALA ?	CEL			32			
ACTINASTRUM	COL		0.3				
ACTINASTRUM HANTZSCHII	FIL						
ANABAENA	FIL			X			
ANABAENA ?	CEL		0.3	32			
ASTERIONELLA FORMOSA	CEL			X			
CARTERIA	CEL						
CERATIUM HIRUNDINELLA	CEL						
F. BRACHYCEAS	CEL						
CHLOROGONIUM	CEL						
CHROOCOCCLUS	CEL						
COCONEIS	COL			X			
COELASTRUM MICROPORUM	COL						
COELASTRUM RETICULATUM	COL						
COSNARIUM	COL						
CRUCIGENIA APICULATA	COL		0.3	32			
CRUCIGENIA TETRAPEDIA	COL		2.4	255			
CRYPTOMONAS	CEL			X			
CYCLOTELLA ATOMUS	CEL						
CYCLOTELLA MENEGHINIANA ?	CEL			X			
CYCLOTELLA MENEGHINIANA ?	CEL		2.7	287			
CYCLOTELLA STELLIGERA	CEL						
CYMBELLA	CEL			X			
CYMBELLA AMPHICEPHALA	CEL						
CACTYLOCCOCCOPSIS	CEL						
DINOBRYON BAVARIUM	CEL						
DINOBRYON DIVERGENS	CEL		0.6	64			
DINOBRYON SOCIALE	CEL		3.0	319			
V. AFRICANUM	CEL						
DINOFAGELLATE	CEL						
DINOFAGELLATE #2	CEL						

LAKE NAPE: BARREN RIVER RES.
STORET NUMBER: 2105

CONTINUED

TAXA	FORM	05 18 73				08 11 73				10 23 73			
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
EUASTRUM	CEL												X
EUGLENA	CEL						0.4					0.4	15
EUGLENA #1	CEL		0.3	32									X
EUGLENA #2	CEL			X									
EUNOTIA	CEL			4402			5.1						938
FLAGELLATES	CEL	11	42.1		4								
GYROSIGMA #1	CEL												
GYROSIGMA #2	CEL												
LYNGBYA LINNETICA	FIL				2	12.2						0.4	15
MELOSIRA #2	CEL	12	12.2	1276								5.4	181
MELOSIRA #4	CEL	15	4.0	415									X
MELOSIRA #5	CEL												
MELOSIRA DISTANS	CEL		3.7	383			0.2					1128.7	968
MELOSIRA VARIANS	CEL			X									
PICRACTINIUM ?	COL			X									
NAVICULA #1	CEL												
NAVICULA ANGLICA													
V. SUBSALSA	CEL												
NAVICULA TRIPUNCTATA	CEL			X									
NITZSCHIA	CEL	4	6.1	638			0.6						X
NITZSCHIA #2	CEL												
NITZSCHIA HOLSATICA ?	CEL												106
OSCILLATORIA	FIL		0.3	32								3.1	
OSCILLATORIA GEMINATA	FIL												
OSCILLATORIA LINNETICA	FIL												
PANDORINA MGRUM	COL						0.4					0.9	30
PEDIASTRUM BIRADIATUM	COL			X									X
PEDIASTRUM SIMPLEX	COL												X
PEDIASTRUM SIMPLEX	COL												X
V. DUODENARIUM	COL												
PENNATE DIATOM	CEL			X									
PENNATE DIATOM #3	CEL											1.8	61

LAKE NAME: BARREN RIVER RES.
STORET NUMBER: 2105

CONTINUED

TAXA	FORM	05 18 73		08 11 73		10 23 73	
		IS	SC	ALGAL UNITS PER ML	IS	SC	ALGAL UNITS PER ML
PHACLS	CEL						
RAPHIDIOPOSTS ?	FIL						
SCENEDESMUS #1	COL		0.3	32		4.5	151
SCENEDESMUS #2	COL			X			X
SCENEDESMUS ABUNCANS	COL						X
SCENEDESMUS DENTICULATUS	CCL					0.4	15
SCENEDESMUS DIMORPHUS	COL						X
SCENEDESMUS QUADRICAUCA	COL			X		2.3	76
STAUROSTRUM	CEL						X
STEPHANODISCUS	CEL						X
SYNEORA ? #1	CEL	3120.1		2105		0.4	15
SYNEORA #1	CEL					1.8	61
SYNEORA DELICATISSIMA	CEL		0.6	64			
SYNEORA DELICATISSIMA ?	CEL		0.6	64		2.7	91
TETRAEDRON	CEL						
TETRAEDRON MINIMUM	CEL						X
TETRAEDRON TRIGONUM	CEL					3.6	121
V. GRACILE	CEL						
TETRASTRUM STAUROGENIAEFORME	CEL						X
TRACHELCOMNAS	CEL			X			
TRACHELCOMNAS #1	CEL						
TRACHELCOMNAS SCHAUINSLANDII	CEL					0.4	15
TRACHELCOMNAS VOLVCCINA	CEL						
TOTAL				10464		14545	3372